

Transitioned Trainees – Paediatric Emergency Requirement (PER)

	Trainees who have logged paediatric cases during advanced training	Trainees who have completed 3 months of paediatric ED during advanced training	Trainees who have not started their paediatric requirement at the time of transition
Required to be completed	Requirement must be completed prior to Fellowship. The PER does not need to be completed by the end of Training Stage 3.		
Number of Paediatric Cases required	Minimum of 400 cases required. Credit is given to cases already completed.	Minimum of 200 cases required. Credit is given to cases already completed outside of the 3-month placement.	Minimum of 400 cases required.
Setting	Future cases must be completed in an accredited Paediatric ED or Mixed ED accredited for the PER.		All cases must be completed in an accredited Paediatric ED or Mixed ED accredited for the PER. Maximum of 200 cases can be completed overseas.
PER-WBAs Required	<p>The number PER-WBAs required is determined by the number of cases left to complete at the time or transition:</p> <p><i>0-49 cases to complete = 1 PER WBA required</i> <i>50-99 cases to complete = 2 PER WBAs required</i> <i>100-149 cases to complete = 3 PER WBAs required</i> <i>150-199 cases to complete = 4 PER WBAs required</i> <i>200-249 cases to complete = 5 PER WBAs required</i> <i>250-299 cases to complete = 6 PER WBAs required</i> <i>300-349 cases to complete = 7 PER WBAs required</i> <i>350-400 cases to complete = 8 PER-WBAs required</i></p> <p><i>The mix of WBA instruments for transition trainees is detailed in Table A below.</i></p>		8 PER-WBAs required (as per Regulation G).
Paediatric Emergency Portfolio	Logbook part only required. DEMT Comment not required.		Required
Age Requirements	No requirements, cases can be up to 15 years, 364 days.		Minimum 200 cases aged <5. Minimum 200 cases aged >5 <15.
Triage requirement	Minimum 100 cases triage 1, 2, 3.		Minimum 150 cases triage 1, 2, 3. Minimum 50 cases must be triage 1, 2 of which at least 25 must be less than five years of age.

TABLE A: Type and number of PER-WBAs required following transition.

No. of PLB cases left to complete	No of PER-WBAs required	PER-WBAs required	PER-WBAs that will be credited
0-49	1	<ul style="list-style-type: none"> 1 PER-DOPS: <u>Procedural Sedation</u> for paediatric patient of five (5) years or less of age 	<ul style="list-style-type: none"> 1 PER-DOPS 3 PER-Mini CEX 3 PER CbD
50-99	2	<ul style="list-style-type: none"> 1 PER-DOPS: <u>Procedural Sedation</u> for paediatric patient of five (5) years or less of age 1 PER-Mini-CEX: Focused assessment of a paediatric patient aged less than two (2) years 	<ul style="list-style-type: none"> 1 PER-DOPS 2 PER-Mini CEX 3 PER CbD
100-149	3	<ul style="list-style-type: none"> 1 PER-DOPS: <u>Procedural Sedation</u> for paediatric patient of five (5) years or less of age 2 PER-Mini-CEX: Focused assessment of a paediatric patient aged less than two (2) years <p>Focused assessment of a paediatric patient aged two (2) to twelve (12) years</p>	<ul style="list-style-type: none"> 1 PER-DOPS 1 PER-Mini CEX 3 PER CbD
150-199	4	<ul style="list-style-type: none"> 1 PER-DOPS: <u>Procedural Sedation</u> for paediatric patient of five (5) years or less of age 3 PER-Mini-CEX: Focused assessment of a paediatric patient aged less than two (2) years <p>Focused assessment of a paediatric patient aged two (2) to twelve (12) years</p> <p>Paediatric patient discharge communication for common diagnosis</p>	<ul style="list-style-type: none"> 1 PER-DOPS 3 PER CbD



200-249	5	<ul style="list-style-type: none"> • 2 PER-DOPS: <u>Procedural Sedation</u> for paediatric patient of five (5) years or less of age <u>Specimen Collection</u> Specimen collection for lab analysis, for a paediatric patient of five (5) years or less of age for any of the following: peripheral intravenous cannula insertion, suprapubic catheter aspiration, in-dwelling urinary catheter aspiration, lumbar puncture • 3 PER-Mini-CEX: Focused assessment of a paediatric patient aged less than two (2) years Focused assessment of a paediatric patient aged two (2) to twelve (12) years Paediatric patient discharge communication for common diagnosis 	<ul style="list-style-type: none"> • 3 PER CbD
300-349	7	<ul style="list-style-type: none"> • 2 PER-DOPS: <u>Procedural Sedation</u> for paediatric patient of five (5) years or less of age <u>Specimen Collection</u> Specimen collection for lab analysis, for a paediatric patient of five (5) years or less of age for any of the following: peripheral intravenous cannula insertion, suprapubic catheter aspiration, in-dwelling urinary catheter aspiration, lumbar puncture • 3 PER-Mini-CEX: Focused assessment of a paediatric patient aged less than two (2) years Focused assessment of a paediatric patient aged two (2) to twelve (12) years Paediatric patient discharge communication for common diagnosis. 	<ul style="list-style-type: none"> • 1 PER CbD



		<ul style="list-style-type: none"> • 2 PER Cbd: One must Less than two (2) years of age • Two (2) to twelve (12) years of age 	
350-400	8	<ul style="list-style-type: none"> • 2 PER-DOPS: <u>Procedural Sedation</u> for paediatric patient of five (5) years or less of age <u>Specimen Collection</u> Specimen collection for lab analysis, for a paediatric patient of five (5) years or less of age for any of the following: peripheral intravenous cannula insertion, suprapubic catheter aspiration, in-dwelling urinary catheter aspiration, lumbar puncture • 3 PER-Mini-CEX: Focused assessment of a paediatric patient aged less than two (2) years Focused assessment of a paediatric patient aged two (2) to twelve (12) years Paediatric patient discharge communication for common diagnosis. • 3 PER Cbd: One must Less than two (2) years of age Two (2) to twelve (12) years of age 1 X Medium complexity 1 X High complexity Required to do all the above 8 PER-WBAs. Further details can be found in the 2022 FACEM Training Program Handbook. 	<ul style="list-style-type: none"> • Nil